The global landscape for NCDs and maternal health

Priya Kanayson, Policy and Advocacy Manager Tuesday 5 November, 2019



The current burden of NCDs




Figure 1: Number of deaths in 2016 from non-communicable diseases (NCDs), injuries, and communicable, maternal, perinatal, and nutritional conditions




NCDs and women

Causes of death for women globally in 2016:"



NCDs caused 19.4 MILLION DEATHS (of 26.5 million), which means



Source: https://www.who.int/healthinfo/global_burden_ disease/estimates/en/

CxCa stat for 2018: https://www.who.int/cancer/cervicalcancer/for-women-and-girls



CVD caused 8.8 MILLION DEATHS among women



861,000 DEATHS were due to DIABETES



1.5 MILLION

women died due to RESPIRATORY CONDITIONS




18-25% of women who live in LMICs are living with DEPRESSION



Over 300,000 women died from CERVICAL CANCER in 2018



NCDs: Common risk factors and solutions



Reduce MODIFIABLE RISK FACTORS



Strengthen HEALTH

SYSTEMS



Increase FUNDING

NCDs at the global level





THANK YOU

SHARE. DISCUSS. ENGAGE. CHANGE.

Ø
f #NCDs
@ncdalliance



MAKING NCD PREVENTION AND CONTROL A PRIORITY, EVERYWHERE